

Child Maltreatment: A Neurodevelopmental Perspective on the Role of Trauma and Neglect in Psychopathology

Bruce D. Perry, M.D., Ph.D

IN: Child and Adolescent Psychopathology

(Edited by Theodore Beauchaine and Stephen P. Hinshaw)

John Wiley & Sons, Hoboken, NJ

2008

pp. 93-129

This is a CTA Web-version of the above chapter

Introduction

The purpose of this chapter is to examine potential mechanisms underlying the well-documented, complex relationships between maltreatment in childhood and the subsequent development of psychopathology. Thousands of studies over the last fifty years have described various aspects of these relationships. Maltreatment in childhood increases risk for virtually every DSM-IV disorder, from autistic-spectrum disorders to schizophrenia to ADHD to major depression to substance abuse disorders. The mechanisms underlying this maltreatment related increase in risk of neuropsychiatric problems are undetermined. The key question addressed in this chapter is “How can abuse lead to psychopathology?” The perspective of the present chapter is neurodevelopmental. This “lens” provides significant insight about the sometimes confusing interrelationships between psychopathology, DSM-IV “diagnoses” and developmental trauma or neglect. A neurodevelopmental perspective is meant to compliment other theoretical and experimental views and can provide useful clues to the mechanisms underlying the origins of neuropsychiatric problems.

The primary premise of a neurodevelopmental perspective is that the human brain is the organ mediating all emotional, social, cognitive and behavioral functioning. Neuropsychiatric disorders and psychopathology, therefore, must involve altered functioning of systems in the brain. The specific nature of dysfunction (e.g., anxiety vs inattention vs affect regulation vs thought disorder) will be determined by which neural networks and brain areas are altered. The present chapter provides an overview of key neurodevelopmental processes and important neural networks which are impacted by abuse and suggests mechanisms which may underlie neuropsychiatric problems related to developmental maltreatment. The major conclusion of this chapter is that we can make plausible conclusions regarding the effects of abuse if we understand how these experiences impact the developing brain. Simply stated childhood trauma will result in alterations in the

systems in the brain which mediate the stress response and neglect will result in dysfunctions in the neural systems which do not receive appropriately timed, patterned repetitive stimulation.

Two major forms of maltreatment will be reviewed in the present chapter: neglect and trauma. Though often co-occurring, these two types of maltreatment are distinctly different in the impact they have on the developing brain, and, therefore, will have differing impacts on the development of psychopathology. Neglect, defined from a neurobiological perspective, is the absence of an experience or pattern of experiences required to express an underlying genetic potential in a key developing neural system. Trauma, from a neurobiological perspective, is an experience or pattern of experiences which activate the stress response systems in such an extreme or prolonged fashion as to cause alterations in the regulation and functioning of these systems. Both neglect- and trauma-related abnormalities in neurodevelopment would be predicted to cause significant psychopathology.

Abuse can have a negative impact on development in several ways. Maltreatment may be the primary mediator of psychopathology when these abnormal experiences directly alter developing neural systems; for example, trauma may cause post-traumatic stress disorder or neglect an attachment disorder. In addition, trauma or neglect may play an exacerbating or expressing role for neuropsychiatric syndromes in individuals with genetic vulnerabilities (e.g., major depression and schizophrenia). And finally, symptoms and problems caused by maltreatment can be disrupting factors for subsequent developmental opportunities (e.g., the disrupting impact of hypervigilance on academic experiences or of neglect-related attachment problems on social development). Often these secondary and tertiary effects are as devastating as the primary abuse-related pathology.

In order to better understand the potential mechanisms by which abuse can cause psychopathology it is crucial to consider the core processes and principles of neurodevelopment

Neurodevelopment

The brain develops rapidly in utero and in the first years of life. During this time important molecular processes are taking place which, if disrupted, can result in abnormal organization and function. Depending upon the nature, timing and frequency of maltreatment, all of these processes can be impacted by chaos, threat, trauma and neglect.

Molecular Processes of Neurodevelopment

Neurogenesis: The vast majority of cell birth or neurogenesis takes place in utero. Few neurons are born after birth, although researchers have demonstrated neurogenesis in the mature brain (Gould, Reeves, Graziano, & Gross, 1999). Neurogenesis may be impacted by maternal substance use, most especially alcohol use and abuse during pregnancy. This can have a devastating impact on the fetal brain growth and result in profound psychopathology later in life (Bookstein et al., 2002).

Migration: As neurons are born and the brain grows, neurons move. It is the fate of some neurons to settle in the brainstem, others in the cortex, for example. Cortical cell

migration and fate mapping are some of the most studied processes in developmental neuroscience (Rakic, 1981; 1996). It is clear that both genetic and environmental factors play important roles in determining a neuron's final location. Migration takes place primarily during the intrauterine and immediate perinatal period but continues throughout childhood and, possibly, to some degree into adult life. A host of intrauterine and perinatal insults – experiences such as infection, lack of oxygen, exposure to alcohol and various psychotropic drugs can alter migration of neurons and have profound impact on the expression of genetic potentials for a host of functions (see Perry, 1988).

Differentiation: Neurons can mature to thousands of unique structures, producing any of a hundred different neurotransmitters (e.g., norepinephrine, dopamine, serotonin, CRF or Substance P). Developing neurons change (differentiate) in response to chemical, often neurochemical, signals. Therefore, any experience that alters neurochemical, hormonal or micro-environmental signals (e.g., extreme activation of the stress-response) during development can change the ways in which certain neurons differentiate, thereby altering the functional capacity of the neural networks in which these neurons reside (e.g., Rutledge et al., 1974).

Apoptosis: More neurons born than are required to make a functional system. Redundant neurons, when unable to adequately “connect” into an active neural network, die (Kuan, Roth, Flavell, & Rakic. 2000). Neurons that make synaptic connections with others and have an adequate level of stimulation will survive; neurons with little activity resorb. This is an example of the important principle of activity-dependent development (see below). Obviously, under stimulation from neglect could increase apoptosis.

Arborization: As neurons differentiate, they send out one form of fiber-like receptive processes called dendrites. The density of these dendritic branches is related to the frequency and intensity of incoming signals. Arborization allows the neuron to receive, process, and integrate complex patterns of input. Dendritic density appears to be one of the most experience-sensitive physical features of a neuron (Diamond, Law, Rhodes, et al. 1966; Greenough, Volkmar, & Juraska. 1973).

Synaptogenesis: The most experience-sensitive feature of a neuron is the synapse. Developing neurons send out fiber-like processes which become axons and synapses. A continuous dynamic of synaptic neurotransmission regulates the activity chains of neurons that allow all brain function. Neurons, during development, find and connect with the appropriate target neurons. This process is guided by certain growth factors and cellular adhesion molecules that attract or repel a specific growth cone to appropriate target neurons. Furthermore this process is influenced by neurotransmission; altering levels of dopamine while nigrostriatal cells are developing will alter receptor expression and synaptogenesis (Wainwright et al., 1995). During the first eight months of life there is an eight-fold increase in synaptic density while the developing neurons in the brain are “seeking” their appropriate connections (Huttenlocher. 1979; 1994). This explosion of synaptogenesis allows the brain to have the flexibility to organize and function with a wide range of potential. Due to the rapid and important neural changes taking place in this first year, this is also a time of remarkable vulnerability to trauma and neglect.

Synaptic sculpting: The synapse is a dynamic structure. With continuous, but episodic release of neurotransmitter, occupation of receptors, release of growth factors, shifts of ions in and out of cells, laying down of new microtubules and other structural molecules, the synapse is continually changing. A key determinant in this synaptic sculpting process is the activity of neurotransmission. When there is a consistent active process of

neurotransmitter release, synaptic connections will be strengthened with actual physical changes that make the pre- and postsynaptic neurons grow closer together, making neurotransmission between these two neurons more efficient. When there is little activity, the synaptic connection will dissolve. Synaptic sculpting is a “use it or lose it” process (see below). Dynamic synaptic sculpting continues through the entire life cycle; however the rate of sculpting decreases with age. This powerful activity-dependent process appears to be the molecular basis of learning, memory and, therefore, at the core of neurodevelopment.

Myelination: Specialized glial cells wrap around axons and, thereby, create more efficient electrochemical transduction down the neuron. This allows a neural network to function more rapidly and efficiently, thereby allowing more complex functioning (e.g., walking depends upon the myelination of neurons in the spinal cord for efficient, smooth regulation of neuromotor functioning.) The process of myelination begins in the first year of life but continues in many key areas throughout childhood with a major burst of myelination in key cortical areas taking place in adolescence. Final myelination of key cortical tracts can take place as late as age 30.

The neurodevelopmental processes described above are sensitive to “signals” from neurotransmitters, neuromodulators, neurohormones, ions, growth factors, cellular adhesion molecules and other morphogens. Disruption of the pattern, timing or intensity of these cues can lead to abnormal neurodevelopment and psychopathology. The specific psychopathology will depend upon the timing of the insult (e.g., was the insult in utero during the development of the brainstem or at age two during the active development of the cortex), the nature of the insult (e.g., is there a lack of sensory stimulation from neglect or an abnormal persisting activation of the stress response from trauma?), the pattern of the insult (i.e., is this a discreet single event, a chronic experience with a chaotic pattern or an episodic event with a regular pattern?). Experience, good and bad, literally becomes the neuroarcheology of the individual’s brain (see Perry, 2001a).

The effects of abuse, then, will be very dependent upon the state of development of the child, and, of course the stage of neurodevelopment. Several key principles help explain the key findings of developmental maltreatment.

Principles of Neurodevelopment

Nature and Nurture: Neurodevelopment is a product of genetic potential and how that potential is expressed as function of the timing, nature and pattern of experience. Genetic differences in key neural factors have been demonstrated that may be related to the functioning of the stress-response systems, for example (Caspi et al., 2002). Yet the way these genetic vulnerabilities are expressed remain sensitive to developmental experiences of trauma (Caspi et al., 2003).

The impact of experience on neurodevelopment shifts during development. In the just fertilized ovum, chemical processes that are driving development are genetically-determined sequences of molecular events. By birth, however, the brain has developed to the point where environmental cues mediated by the senses play a major role in neural differentiation, arborization, and synaptogenesis, helping create functional neural networks. By adolescence, the majority of the changes that are taking place in the brain of that child are determined by experience, not genetics. The languages, beliefs, cultural practices, and

complex cognitive and emotional functioning (e.g., self esteem) by this age are primarily experience-based.

Sequential Developmental: The brain develops in a sequential and hierarchical fashion; organizing itself from least (brainstem) to most complex (cortical areas). These different areas develop, organize and become fully functional at different times during childhood. At birth, for example, the brainstem areas responsible for regulating cardiovascular and respiratory function must be intact for the infant to survive, and any malfunction is immediately observable. In contrast, the cortical areas responsible for abstract cognition are not fully functional until adult life.

This is important in understanding the role of the stress-response neural systems in shaping future neurodevelopment. As described in detail below, the monoamine neurons originating in the brainstem and diencephalon (present by birth) send direct connections into all other brain areas (Figure 1). If these systems are poorly organized or dysregulated themselves, they can dysregulate and disorganize higher parts of the brain. The “higher” more complex areas (e.g., limbic and cortical) have not yet organized. The final organization of these important areas (and all of the functions mediated by these areas) will be influenced by the pattern, intensity and nature of stimulation coming from these adrenergic, noradrenergic, serotonergic and dopaminergic neurons.

Trauma will result in profound alterations in the pattern of activation in these stress-mediating neural systems; which in turn will result in patterned, repetitive neuronal activation in a distributed and diverse set of brain systems, influencing the organization and function of these higher brain areas; the result can be compromised function and psychopathology. Traumatic stress, for example, can impact cortically-mediated (e.g., cognition), limbic-mediated (e.g., affect regulation), diencephalic-mediated (e.g., fine motor regulation, startle response) and brainstem-mediated (e.g. heart rate, blood pressure regulation) functioning.

Each brain area will have its own timetable for development. The neurodevelopmental processes described above will be most active in different brain areas at different times and will, therefore, either require (critical periods) or be sensitive to (sensitive periods) organizing experiences (and the neurotrophic cues related to these experiences). The neurons for the brainstem have to migrate, differentiate and connect, for example, before the neurons for the cortex.

Disruptions of experience-dependent neurochemical signals during early life may lead to major abnormalities or deficits in neurodevelopment. Disruption of critical neurodevelopmental cues can result from 1) lack of sensory experience during sensitive periods (e.g., neglect) or 2) atypical or abnormal patterns of necessary cues due to extremes of experience (e.g., traumatic stress, see Perry, 2001a; 2001b). This has implications for understanding psychopathology related to maltreatment. In the development of socio-emotional functioning, early life nurturing appears to be critical, for example. If this is absent for the first three years of life and then a child is adopted and begins to receive attention, love and nurturing, these positive experiences may not be sufficient to overcome the malorganization of the neural systems mediating socio-emotional functioning.

Activity-dependent Neurodevelopment: The brain organizes in a use-dependent fashion. Many of the molecular processes in neurodevelopment are activity dependent. In the developing brain, undifferentiated neural systems are critically dependent upon sets of

environmental and micro-environmental cues (e.g., neurotransmitters, cellular adhesion molecules, neurohormones, amino acids, ions) in order for them to appropriately organize from their undifferentiated, immature forms (Lauder, 1988; Perry, 1994) (Perry & Pollard, 1998). Lack, or disruption, of these critical cues can alter the neurodevelopmental processes of neurogenesis, migration, differentiation, synaptogenesis - all of which can contribute to malorganization and diminished functional capabilities in the specific neural system where development has been disrupted. This is the core of a neuroarcheological perspective on dysfunction related adverse childhood events (Perry, 2001b).

These molecular cues that guide development are dependent upon the experiences of the developing child. The quantity, pattern of activity and nature of these neurochemical and neurotrophic factors depends upon the presence and the nature of the total sensory experience of the child. When the child has adverse experiences - loss, threat, neglect, and injury - there can be disruptions of neurodevelopment that will result in neural organization that can lead to compromised functioning throughout life (see below). Healthy organization of neural networks depends upon the pattern, frequency and timing of key experiences during development. Patterned, repetitive activity changes the brain. Chaotic, episodic experiences out of sync with the developmental stage of the child create chaotic, developmentally-delayed dysfunctional organization.

If a child is neglected - if he or she hears fewer words, has fewer relational opportunities, has less physical comfort and love - the rapidly organizing networks in the brain which mediate language, social affiliation and attachment will not receive sufficient patterned, repetitive activation to develop normally. The result is a neglect related set of deficits. The deficit will be in the domains where the neglect occurred (see Perry 2002).

Sensitive and Critical Windows: Due to sequential and activity-dependent development of the brain there are times when a given developing neural system is more sensitive to experience. In healthy development, that sensitivity allows the brain to efficiently organize in response to the unique demands of a given environment. If the child's is born into a hunter-gatherer clan in New Guinea as opposed to a suburban English family, for example, different genes can be expressed, and different neural networks can be organized to best fit that family, culture and environment. This sensitivity, for most functions, passes with time. Acquisition of language with full fluency after a certain age becomes more difficult; formation of key relational and attachment capacities after the first five years of life becomes difficult if these initial years were characterized by disorganized, absent or abusive primary caregivers.

Sensitive periods are different for each brain area and neural system, and therefore, for different functions. The sequential development of the brain and the sequential unfolding of the genetic map for development mean that the sensitive periods for neural system (and the functions they mediate) will be when that area is most actively organizing. The brainstem must organize key systems by birth; therefore, the sensitive period for those brainstem-mediated functions is during the prenatal period. A variety of in utero experiences may influence the development of the key stress-mediating neurotransmitter systems which originate in the brainstem. For example, prenatal exposure to psychoactive drugs may disrupt normal development of the brainstem catecholamines (Perry, 1988). In animal models, brief prenatal and perinatal stress can cause altered development of hippocampal organization and the hypothalamic-pituitary-adrenal axis which remain altered throughout life (Plotsky and Meaney, 1993; Shors et al., 1990). The neocortex, in contrast,

has systems and functions organizing throughout childhood and into adult life. The sensitive periods for these cortically mediated functions are likely to be very long.

The primary clinical implication of this is that early childhood trauma or maltreatment has disproportionate capacity to cause significant dysfunction in comparison with similar trauma or neglect later in life (see Rutter et al., 1998; Rutter et al., 1999). Children are more vulnerable to trauma and neglect than adolescents and adults. The younger a child is, the more likely they are to have enduring and pervasive problems following trauma and neglect. Neglect in the first years of life can have devastating impact even if a child is removed from the neglectful environment (Perry 2002; Perry, 2005). The longer a child remains in a neglectful environment, the more vulnerable they become (e.g., Rutter et al. 1998; 1999; O'Connor et al. 2000; Perry, 2005).

The major consequence of these principles of brain development is that the organizing brain of an infant or young child is more malleable to experience than a mature brain. While experience may alter the behavior of an adult, experience literally provides the organizing framework for an infant and child. Because the brain is most plastic (receptive to environmental input) in early childhood, the child is most sensitive to experience at this time – both good and bad.

The Neurodevelopmental Impact of Neglect

Deprivation during the sensitive period of a given neural system will result in insufficient patterned, repetitive activity to stimulate adequate organization. The deprived developing neural network will have altered neural microarchitecture which may include cell migration, synaptogenesis, dendritic sprouting. The result is a neural system less functionally capable. This deprivation – neglect – can take multiple forms. In some rare cases, a single domain of functioning is primarily impacted. Our clinical group has worked with several children raised with cognitive stimulation, physical and emotional warmth but physically restrained for multiple years and, therefore, unable to sit, stand or walk. Most neglect takes the form of chaotic, mis-timed, inconsistent experience related to the isolation, personal chaos, incompetence, ignorance, domestic violence, substance abuse or psychopathology of the primary caregiver. This can manifest as delays in motor, self-regulatory, affective and cognitive functioning. A common manifestation of this form of neglect is in speech and language acquisition; if the infant or toddler hears few words, the developing speech and language systems will be impacted (Huttenlocher et al., 2002); the result is impaired speech and language.

The earlier and more pervasive the neglect is, the more devastating the developmental problems for the child. Indeed, a chaotic, inattentive and ignorant caregiver can produce pervasive developmental delay (PDD; DSM IV-R) in a young child (Rutter, Andersen-Wood, Beckett, et al. 1999). Yet the very same inattention for the same duration if the child is ten will have very different and less severe impact than inattention during the first years of life.

Studies of the neurodevelopmental impact of neglect are not as common as those on the effects of trauma. Both animal studies, descriptive reports with severely neglected children and some recent studies with adopted children following neglect which document some aspects of the neurodevelopmental impact of neglect.

Animal Studies: Hubel and Weisel's landmark studies on development of the visual system using sensory deprivation techniques helped define the concepts of critical and sensitive period (Hubel & Wiesel. 1963). In hundreds of other studies, extremes of sensory deprivation (Hubel & Wiesel. 1970; Greenough, Volkmar, & Juraska. 1973) or sensory enrichment (Greenough & Volkmar. 1973; Diamond, Krech, & Rosenzweig. 1964; Diamond, Law, Rhodes, et al. 1966) have been studied. These include disruptions of visual stimuli (Coleman & Riesen. 1968), environmental enrichment (Altman & Das. 1964; Cummins & Livesey. 1979), touch (Ebinger. 1974; Rutledge, Wright, & Duncan. 1974), and other factors that alter the typical experiences of development (Uno, Tarara, Else, & et.al. 1989; Plotsky & Meaney. 1993; Meaney, Aitken, van Berkal, Bhatnagar, & Sapolsky. 1988).

These findings generally demonstrate that the brains of animals reared in enriched environments are larger, more complex and functional more flexible than those raised under deprivation conditions. Relationships between experience and brain cytoarchitecture have demonstrated a relationship between density of dendritic branching and the complexity of an environment (Diamond & Hopson. 1998). Rats raised in environmentally enriched environments have higher density of various neuronal and glial microstructures, including a 30% higher synaptic density in cortex compared to rats raised in an environmentally deprived setting (Bennett, Diamond, Krech, & Rosenzweig. 1964; Altman & Das. 1964). Animals raised in the wild have from 15 to 30% larger brain mass than their offspring who are domestically reared (Darwin. 1868; Rohrs. 1955; Rohrs & Ebinger. 1978; Rehkamper, Haase, & Frahm. 1988).

Animal studies suggest that critical periods exist during which specific sensory experience was required for optimal organization and development of the part of the brain mediating a specific function (e.g., visual input during the development of the visual cortex). While these phenomena have been examined in great detail for the primary sensory modalities in animals, few studies have examined the issues of critical or sensitive periods in humans. What evidence there is would suggest that humans tend to have longer periods of sensitivity and that the concept of critical period may not be useful in humans (Perry, 2006). Yet altered emotional, behavioral, cognitive, social and physical functioning has been demonstrated in humans following specific types of neglect. For obvious ethical reasons, most of these findings come from descriptive clinical rather than experimental studies.

Neglect in Early Childhood: The majority of clinical reports on neglect have focused on institutionalized children or feral children. As early as 1833, with the famous Kaspar Hauser, feral children had been described (Heidenreich. 1834). Hauser was abandoned as a young child and raised from early childhood (likely around age two) until seventeen in a dungeon, experiencing relative sensory, emotional and cognitive neglect. His emotional, behavioral and cognitive functioning was, as one might expect, very primitive and delayed.

In the early forties, Spitz described the impact of neglectful caregiving on children in foundling homes (orphanages). Most significant, he was able to demonstrate that children raised in fostered placements with more attentive and nurturing caregiving had superior physical, emotional and cognitive outcomes (Spitz. 1945; Spitz. 1946).

Dennis (1973) described a series of findings from children raised in a Lebanese orphanage. These children were raised in an institutional environment devoid of individual attention, cognitive stimulation, emotional affection or other enrichment. Prior to 1956 all of these children remained at the orphanage until age six, at which time they were

transferred to another institution. Evaluation of these children at age 16 demonstrated a mean IQ of approximately 50. When adoption became common, children adopted prior to age 2 had a mean IQ of 100 by adolescence while children adopted between ages 2 and 6 had IQ values of approximately 80 (Dennis, 1973). This graded recovery is consistent with the known principles of neurodevelopment described above; the older a child was at time of adoption, (i.e., the longer the child spent in the neglectful environment) the more pervasive and resistant to recovery were the deficits.

Money and Anncillo (1976) reported the impact of change in placement on children with psychosocial dwarfism (failure to thrive). In this preliminary study, 12 of 16 children removed from neglectful homes recorded remarkable increases in IQ and other aspects of emotional and behavioral functioning. Furthermore, they reported that the longer the child was out of the abusive home the higher the increase in IQ. In some cases IQ increased by 55 points.

A more recent report on a group of 111 Romanian orphans (Rutter & English and Romanian Adoptees study team, 1998; Rutter, Andersen-Wood, Beckett, et al. 1999) adopted prior to age two from very emotionally and physically depriving institutional settings demonstrate similar findings. Approximately one half of the children were adopted prior to age six months and the other half between six months and 2 years old. At the time of adoption, these children had significant delays. Four years after being placed in stable and enriching environments, these children were re-evaluated. While both groups improved, the group adopted at a younger age had a significantly greater improvement in all domains. As a group, these children were at much greater risk for meeting diagnostic criterion for autism-spectrum disorder, a finding that sheds light on the evolving relationships between early life trauma, neglect and subsequent development of severe neuropsychiatric problems including psychotic disorders and schizophrenia (Read et al., 2001).

These observations are consistent with the clinical experiences of the ChildTrauma Academy working with maltreated children for the last fifteen years. During this time we have worked with more than 1000 seriously children. We have recorded increases in IQ of over 40 points in more than 60 children following removal from neglectful environments and placed in consistent, predictable, nurturing, safe and enriching placements (Perry, 2005). In a group of 200 children under the age of 6 at time of removal, significant developmental delays were seen in more than 85% of the children. The severity of these developmental problems increased with age, suggesting, again, that the longer the child was in the neglectful environment - the earlier and more pervasive the neglect - the more indelible and pervasive the deficits.

Neurobiological Findings

All of these reported developmental problems – language, fine and large motor delays, impulsivity, disorganized attachment, dysphoria, attention and hyperactivity, and a host of others described in these neglected children – are caused by abnormalities in the brain. Despite this obvious statement, very few studies have examined directly any aspect of neurobiology in neglected children. One early clue in humans On autopsy, the brain of Kasper Hauser (see above) was notable for small cortical size and few, non-distinct cortical gyri – all consistent with cortical atrophy (Simon, 1978).

Our group has examined various aspects of neurodevelopment in neglected children ((Perry & Pollard, 1997; Perry, 2005). Globally neglected children had smaller the frontal-occipital circumference, a measure of head size and in young children a reasonable

measure of brain size. Neuroimaging demonstrated that 64.7 % of the brain scans were abnormal in the children with global neglect and 12 % in children following chaotic neglect. The majority of the findings were "enlarged ventricles" or "cortical atrophy" (Figure 2). Once removed from the neglectful settings, recovery of function and relative brain-size were observed. The degree of recovery over a year period however was inversely proportional to age in which the child was removed from the neglecting caregivers. The earlier in life and the less time in the sensory-depriving environment, the more robust the recovery (Perry, 2005).

In the study of Romanian orphans described above, the 38 % had FOC values below the third percentile (greater than 2 SD from the norm) at the time of adoption. In the group adopted after six months, fewer than 3 % and the group adopted after six months 13 % had persistently low FOCs four years later (Rutter & English and Romanian Adoptees study team. 1998; O'Connor, Rutter, & English and Romanian Adoptees study team. 2000). Strathearn (2001) has followed extremely low birth weight infants and shown that when these infants end up in neglectful homes they have a significantly smaller head circumference at 2 and 4 years, but not at birth. This is despite having no significant difference in other growth parameters.

Studies from other groups report similar altered neurodevelopment in neglected children. Teicher has reported altered corpus callosum development in neglected children (Teicher et al., 2004). Altered brain-related measures (e.g., salivary cortisol) have been demonstrated in children adopted following neglect (Gunnar et al., 2001). Chugani and colleagues have demonstrated decreased metabolic activity in the orbital frontal gyrus, the infralimbic prefrontal cortex, the amygdale and head of the hippocampus, the lateral temporal cortex and in the brainstem in a group of Romanian orphans (Chugani, et al., 2001).

Neurodevelopmental Impact of Trauma

The brain's stress mediating systems are widely distributed; involve brain and the autonomic nervous system as well as neuroendocrine and neuroimmune responses. Clearly stress-related neural networks permeate the entire brain. The capacity, then, of a use-dependent alteration in the organization and functioning of neural systems involved in dozens of functions could easily take place with prolonged activation. The specific nature of which of these myriad systems becomes altered depends upon the specific nature of the threat response in any given circumstance.

The human brain is continually sensing, processing, storing, perceiving and acting in response to information from the external and internal environments. This continuous monitoring process is especially sensitive to input that may indicate threat. These "worlds" – external as 'sensed' by our five senses and internal as 'sensed' by a set of specialized neurons throughout the body (e.g., glucose or sodium sensitive neurons) – are always changing. Our physiology and neurophysiology are characterized by a continuous process of modulation, regulation, compensation, activation – all designed to keep our body's systems in some state of equilibrium or homeostasis. Whenever the incoming information from either inside or outside the body alter this homeostasis (Perry and Pollard,) or indicate similarity to a pattern of activity previously associated with a threat (Perry,),

the brain initiates compensatory, adaptive responses to re-establish homeostasis or to take the necessary actions to survive.

Sequential Processing of Threat Related Neural Activity

This process begins when our senses transform forms of energy (e.g., light, sound, pressure) into patterned activity of sensory neurons. The first 'stops' for sensory input from the outside environment (e.g., light, sound, taste, touch, smell) and from inside the body (e.g., glucose levels, temperature) are the lower, 'regulatory' areas of the brain (see later). These neural patterns of activity created by sensory input first come into the brain separately – visual input comes into one nuclei, auditory another, olfactory another and so on. The “first stops” for primary sensory input are in lower parts of the brain that are incapable of conscious perception.

Brainstem and Diencephalon

Neural input from our senses directly connects to the lower areas of the CNS in the brainstem, diencephalon and hypothalamus. For example, our internal organs directly relay information to the amygdala and locus coeruleus directly or through the nucleus paragigantocellularis and nucleus tractus solitaries (Elam et al., 1986; Nauta and Whitlock, 1956; Saper, 1982). Other primary sensory input from visual, auditory, tactile and olfactory systems directly connect into these lower brain nuclei where the process of sorting, integrating, interpreting, storing (if appropriate) and responding to these incoming signals begins.

These brainstem and diencephalic nuclei project to the thalamus which begins the process of integrating this information, relaying sensory information to the primary sensory receptive areas of the cortex. These primary sensory regions project to adjacent cortical association areas. A feedback process involves projections from key cortical areas back to the lower parts of the brain; visual, auditory and somatosensory cortical association areas send projections to the hippocampus, amygdala, orbitofrontal cortex, entorhinal cortex, and cingulate gyrus and other brain structures. This reciprocal processing allows the brain to sort, process and “act” on the threat related signals from the body and the external world.

The Reticular Activating System: Key to this entire process is the role of an array of important neurotransmitter networks; the monoamine systems; epinephrine, norepinephrine, dopamine and serotonin – have key nuclei (clusters of cell bodies) which send direct axonal projections to virtually all other areas of the brain and lower to influence the autonomic neurons which leave the brain and directly influence heart, lung, gut, skin and the rest of the organs of the body. Collectively this network has been referred to as the reticular activating system.

The reticular activating system (RAS) is a network of ascending, arousal-related neural systems in the brain that consists of locus ceruleus noradrenergic neurons, dorsal raphe serotonergic neurons, cholinergic neurons from the lateral dorsal tegmentum, and mesolimbic and mesocortical dopaminergic neurons, among others. Much of the original research on arousal, fear, and response to stress and threat was conducted using various lesion models of the reticular activating system (Moore and Bloom, 1979). The reticular activating system appears to be an integrated neurophysiological system involved in arousal, anxiety, and modulation of limbic and cortical processing (Munk et al., 1996). Working together, the brain-stem monoamine systems in the reticular activating system

provide the flexible and diverse functions necessary to modulate stress, distress and trauma. A key component of the RAS is the locus coeruleus.

The Locus Coeruleus: The locus coeruleus is involved in initiating, maintaining, and mobilizing the total body response to threat (Aston-Jones et al., 1986). A bilateral grouping of norepinephrine-containing neurons originating in the pons, the locus coeruleus sends diverse axonal projections to virtually all major brain regions and thus functions as a general regulator of noradrenergic tone and activity (Foote et al., 1983). The locus coeruleus plays a major role in determining the “valence,” or value, of incoming sensory information; in response to novel or potentially threatening information, it increases its activity (Abercrombie and Jacobs, 1987a; Abercrombie and Jacobs, 1987b). The ventral tegmental nucleus also plays a part in regulating the sympathetic nuclei in the pons/medulla (Moore and Bloom, 1979). Acute stress results in an increase in locus coeruleus and ventral tegmental nucleus activity and the release of catecholamines throughout the brain and the rest of the body. These brain-stem catecholamine systems (locus coeruleus and ventral tegmental nucleus) play a critical role in regulating arousal, vigilance, affect, behavioral irritability, locomotion, attention, and sleep, as well as the startle response and the response to stress (Levine et al., 1990; Morilak et al., 1987a; Morilak et al., 1987b; Morilak et al., 1987c).

Hypothalamus and Thalamus: Sensory thalamic areas receive input from various afferent sensory systems, and at this level, “feeling” begins. Although thalamic nuclei are important in the stress response, these regions have been studied primarily as “waystations” that transmit important arousal information from the reticular activating system neurons (e.g., locus coeruleus noradrenergic neurons) to key limbic, subcortical, and cortical areas involved in sensory integration and perception of threat-related information (Castro-Alamancos and Connors, 1996). The neuroendocrinological—and likely neuroimmunological—afferent and efferent wings of the threat response are mediated by hypothalamic and other anatomically related nuclei. Studies have demonstrated important roles for various hypothalamic nuclei and hypothalamic neuropeptides in the stress response in animals (Bartanusz et al., 1993; Miaskowski et al., 1988) (Rosenbaum et al., 1988), and humans (Young and Lightman, 1992).

The Limbic System

The central role of the subcortical network of brain structures in emotion was hypothesized by Papez (Papez, 1937) and elaborated by MacLean who coined the term limbic system. This sub-cortical network is responsible for a range of emotion- and relational-related functions. Among the key subcomponents of the limbic system are two brain areas known to be intimately involved in the stress response: the amygdala and hippocampus.

Amygdala: The amygdala has emerged as the key brain region responsible for the processing, interpretation, and integration of emotional functioning (Clugnet and LeDoux, 1990). Just as the locus coeruleus plays the central role in orchestrating arousal, the amygdala plays the central role in the brain in processing afferent and efferent connections related to emotional functioning (LeDoux et al., 1988; Pavlides et al., 1993b; Phillips and LeDoux, 1992b). The amygdala receives input directly from the sensory thalamus, the hippocampus (via multiple projections), the entorhinal cortex, and the sensory association

and polymodal sensory association areas of the cortex as well as from various brain-stem arousal systems via the reticular activating system(Selden et al., 1991). The amygdala processes and determines the emotional valence of simple sensory input, complex multisensory perceptions, and complex cognitive abstractions, even responding specifically to complex socially relevant stimuli. In turn, the amygdala orchestrates the organism's response to this emotional information by sending projections to brain areas involved in motor (behavioral), autonomic nervous system, and neuroendocrine areas of the CNS (Davis, 1992a; Davis, 1992b; LeDoux et al., 1988). In a series of landmark studies, LeDoux and colleagues demonstrated the key role of the amygdala in "emotional" memory (LeDoux et al., 1990). Animals, including humans, store emotional as well as cognitive information, and the storage of emotional information is critically important in both normal and abnormal regulation of anxiety. The "site" at which anxiety is perceived is the amygdala (Davis, 1992a). It is in these limbic areas that the patterns of neuronal activity associated with threat-and mediated by the monoamine neurotransmitter systems of the reticular activating system—become an "fear."

Hippocampus: This brain area is involved in the storage of various kinds of sensory information and is very sensitive to "stress" activation (Pavlides et al., 1993a; Phillips and LeDoux, 1992a; Sapolsky et al., 1984). The hippocampus appears to be critical in the storage and recall of cognitive and emotional memory(Selden et al., 1991).Any emotional state related to arousal or threat may alter hippocampal functioning, changing the efficiency and nature of hippocampal storage and retrieval. Threat alters the ability of the hippocampus and connected cortical areas to "store" certain types of cognitive information (e.g., verbal) but does not affect the storage of other types (e.g., nonverbal).

Hormonal signals affect heterogeneous corticosteroid nuclear receptors, i.e. type 1 (mineralocorticoid) or type 2 (glucocorticoid) in the hypothalamic-pituitary-adrenal (HPA) axis. Stressful life events such as isolation increase HPA axis activity(McEwen, 2001). The hippocampus, amygdala, and mPFC are limbic structures that are targets for and also modulate adrenal steroids. Glucocorticoids can result in neurotoxic damage to the hippocampus with suppression of neurogenesis (McEwen, 2001; Sapolsky, 2000). Exposure to stress results in release of corticotrophin releasing hormone (CRH), adrenocorticotrophic hormone (ACTH), and cortisol via activation of the HPA axis. During periods of stress there is partial resistance to feedback inhibition of cortisol release and increase in plasma cortisol levels, in addition to a decrease in glucocorticoid receptors(Sapolsky and Plotsky, 1990). Glucocorticoid receptors are present in the brain in high density in areas relevant to stress and anxiety such as the hypothalamus, hippocampus, serotonergic and noradrenergic cell bodies on both neurons and glia. Based on animal studies, mineralocorticoid expression is high in limbic regions such as hippocampus, septum and amygdala(Reul and de Kloet, 1985; Veldhuis and De Kloet, 1982). Animal studies suggest that stress experienced during critical years of development can have long lasting effects on HPA axis. For instance, rats that experience in utero stress or early maternal deprivation have increased corticosterone concentrations when exposed to stress. Early postnatal stress is associated with changes in basal concentrations of hypothalamic CRH, mRNA, hippocampal glucocorticoid receptor mRNA, and median eminence CRH, in addition to the stress-induced CRH, cortocosterone, and ACTH release(Levine et al., 1993a; Levine et al., 1993b; Stanton et al., 1988). Adults with PTSD and non-human primates with early adverse experiences have elevated CRH concentrations and decreased cortisol levels in the cerebrospinal fluid (Coplan et al., 1996). Finally a number of studies are indicating the crucial role of corticotrophin-releasing factor

(CRF) and the sensitivity of CRF receptors in mediating stress reactivity in humans (Kehne, 2007).

Cortex

The quality and intensity of any emotional response, including anxiety, are dependent on subjective interpretation or cognitive appraisal of the specific situation eliciting the response (Maunsell, 1995; Singer, 1995). Most theories addressing the etiology of anxiety disorders focus on the process by which stimuli are “mis-labeled” as being “threat” related, thereby inducing a fear response and anxiety in situations where no true threat exists. How individuals “cortically interpret” the limbic-mediated activity (i.e., their internal state) associated with arousal plays a major role in their subjective sense of anxiety (Gorman et al., 1989). Klüver-Bucy syndrome, which results from damage to or surgical ablation of the temporal lobes, is characterized by absence of fear in response to current and previously threatening cues (Klüver and Bucy, 1937). The general disinhibition characteristic of this syndrome suggests loss of the capacity to interpret incoming threat-related cues from lower brain areas.

Heterogeneity of Adaptive Responses to Threat: Hyperarousal and Dissociation

Individual responses to threat can vary tremendously. This is not surprising considering the vast distribution of neural functions which are available to the stress-response network. This network involves the entire brain, and, indirectly, the whole body. This allows the response to potential threat to be appropriate and proportional to the need. The specific adaptive changes taken by the brain to respond to the incoming threat-related signals will vary depending upon many factors; different elements of the widely distributed neural system will be recruited and others will be shut down to conserve energy and focus the body's response to threat. Under normal circumstances (i.e., a normal stress-response capability), the responses are graded, proportional to the level of perceived threat; when the threat is mild, a moderate activation of key systems takes place; when extreme, intense and prolonged activation will occur. Further, adaptive responses to threat are specific to the nature of the threat; either preparing to flee or fight or preparing to be overwhelmed and injured. In cases of abnormal development or sensitivity of the stress-response systems (see later sections) the responses to potential threat are inappropriate and out of proportion; trauma can make the system over-active and overly reactive.

Two major, inter-related response patterns – hyperarousal and dissociative - have been described (Perry et al., 1995). The hyperarousal response has been well characterized; it was originally described as the fight or flight response (Cannon, 1914). As described above, incoming signals activate the locus coeruleus and through a cascade of neural activation recruiting key limbic and cortical areas to focus on, and respond to, the threat. These neural and neuroendocrine activations prepare the body to fight or flee. Cortisol and adrenaline course through the body; heart rate increases, glycogen is mobilized from muscles; all distracting information is tuned out.

However, when fighting or fleeing is not possible, the brain will recruit a different set of neural systems and utilize avoidant, dissociative adaptations. Dissociation is basically a mental mechanism by which one withdraws attention from the outside world and focuses on

the inner world. Dissociation may involve a distorted sense of time, a detached feeling that you are “observing” something happen to you as if it is unreal, the sense that you may be watching a movie of your life. In extreme cases, especially if the trauma is repetitive and painful (e.g., sexual abuse), the child may withdraw into an elaborate fantasy world where she may assume special powers or strengths. Like the alarm response, this “defeat” or dissociative response is graded. The intensity of the dissociation varies with the intensity and duration of the traumatic event.

The neurobiology of dissociation is related but somewhat different from that of the hyperarousal response. Both utilize the monoamine systems in the brainstem and diencephalon; but somewhat different elements of these complex networks. In animals, the ‘defeat’ response has a distinct neurobiology which is similar to dissociation response in humans. Indeed, the neurobiology and phenomenology of dissociation appears to most approximate the ‘defeat’ reaction described in animals (Henry, Stephens, and Ely, 1986; Heinsbroek, van Haaren, Feenstra, and Boon, 1991; Miczek et al., 1990). As with the hyperarousal/fight or flight response, dissociation involves brainstem-mediated CNS activation which results in increases in circulating epinephrine and associated stress steroids (Glavin, 1985; Henry, Liu, Nadra, Qian, Mormede, Lemaire, Ely, and Hendley, 1993; Herman, Guillonneau, Dantzer, Scatton, Semerdjian-Rouquier, and Le Moal., 1982). A major CNS difference, however, is that, in dissociation, vagal tone increases dramatically, decreasing blood pressure and heart rate (occasionally resulting in fainting) despite increases in circulating epinephrine. In addition, there appears to be an increased relative importance of dopaminergic systems, primarily mesolimbic and mesocortical (Kalivas, 1985; Kalivas, Richardson-Carlson, and Van Orden, 1986; Kalivas, Duffy, Dilts and Abhold, 1988; Abercrombie, Keefe, DiFrischia, and Zigmond 1989). These dopaminergic systems are intimately involved in the reward systems, affect modulation (e.g., cocaine-induced euphoria) and, in some cases, are co-localized with endogenous opioids mediating pain and other sensory processing. These opioid systems are clearly involved in altering perception of: painful stimuli, sense of time, place and reality. Indeed, most opiate agonists can induce dissociative responses. Of primary importance in mediating the freeze or surrender dissociative response are endogenous opioid systems (Abercrombie and Jacobs, 1988).

For most children and adults, however, the adaptive response to an acute trauma involves a mixture of hyperarousal and dissociation. During the actual trauma, a child will feel threatened and the arousal systems will activate. With increased threat, the child moves along the arousal continuum. At some point along this continuum, the dissociative response is activated. This results in the host of protective mental (e.g., decreases in the perception of anxiety and pain) and physiological responses (decreased heart rate) that characterize the dissociative response.

Whatever the adaptive response during a trauma, the key issue for subsequent psychopathology is how long these systems are activated. The longer and more intense the activation during the actual traumatic event (s), the more likely there will be molecular changes in the stress-response systems that lead to long-term functional changes. Trauma can cause alterations that lead to sensitized, dysfunctional neural networks; essentially the state of fear can become the persisting trait of anxiety. What were once adaptive neurobiological states can become, over time, maladaptive traits (Perry et al., 1995)

Trauma Alters Stress-Mediating Neural Networks

The clinical impact of traumatic stress on the developing child has been well documented. The simplest are the studies examining the development of obvious trauma-related psychopathology such as post-traumatic stress disorder (for review see; Perry, 1994; 2001b; Glaser, 2000; Teicher et al., 2002; DeBellis and Thomas, 2003; Bremner, 2003). The increased incidence of PTSD following trauma, the list of attenuating and exacerbating has been well documented. Table 1 summarizes the key factors which appear to be related to subsequent development of trauma-specific psychopathology.

Traumatic stress results in altered measures of brain function and in brain-mediated functioning in children. These include measures of hippocampal function, adrenergic receptor functioning, hippocampal and cortical structural development, cardiovascular functioning and emotional, social and behavioral functioning (Perry, 1998; Teicher et al., 1994; 1997; De Bellis et al., 1994; 1997; 1999a; 1999b, 1997; Scaer et al, 2001; Carrion et al., 2001; 2002a). Magnetic resonance imaging (MRI) has revealed reductions in hippocampus (Bremner et al., 1997; Stein et al., 1997; Bremner et al., 2003a), alterations in cerebellar vermis (Anderson, et al., 2002) and altered amygdala (Driessen et al., 2000; Schmahl et al., 2003) volumes as well as deficits in verbal declarative memory measured with neuropsychological testing among women who were sexually abused as children. Sexually abused girls demonstrate neuroendocrine abnormalities as adolescents (Putnam, 1998).

Functioning of monoamine systems in adults is influenced by developmental trauma (Perry, 1995; 2001a; 2001b). Further, developmental trauma appears to influence genetic expression of at least one potential genetic marker for depression. A study of a polymorphism for the promoter region of the serotonin transporter (5-HTT) gene found that childhood maltreatment increased the risk of depression in early adulthood for persons with the common "short" allele compared to persons with the long allele; the short allele is associated with lower transcriptional efficiency of the promoter (Caspi, 2003).

The most overwhelming evidence for the impact of developmental trauma on stress-related neural networks and their functioning comes from a retrospective epidemiological study of 17,000 adults. Over the last ten years the Adverse Childhood Experience (ACE) studies, have been reporting on increased risk of a host of emotional, social, behavioral and physical health problems following abuse and related traumatic experiences in childhood (Anda et al.1991; 2001; Dube et al., 2001a; 2001b; 2002a; 2002b; 2003a; 2003b; . These epidemiological findings converge with evidence from neurobiology about numerous effects of childhood stress on brain and physical systems (Glaser, 2000). These epidemiological studies examined the relationship between adverse childhood experiences including child abuse and a wide range of functioning in adult life. The findings are consistent with the view that developmental trauma impacts the stress-response systems, and, therefore, can have destructive impact on all of the neural systems and functions that are interconnected to this widely distributed network (see Fig. 1).

Among the ACE findings are a graded increase in risk (i.e., more abuse = more risk) for affective symptoms and panic attacks; for memory problems; for hallucinations; for poor anger control; for perpetrating partner violence; unhealthy sexual behavior (early intercourse, promiscuity, sexual dissatisfaction); suicide; substance abuse; alcohol use and abuse; smoking. In addition there is a significant increased risk for a range of physical health problems following child abuse. Risk for the major causes of death in adult life is increased following adverse childhood experiences (Felitti et al.,1998).

Taken together there is little doubt that developmental trauma alters key neural systems involved in mediating the stress response, and, thereby, results in a host of neuropsychiatric and related functional problems.

Summary

All of the major molecular processes involved in brain development can be impacted by abuse. With either neglect or trauma, the timing (the earlier in life the more impact), intensity, pattern and duration of the maltreatment can alter virtually every brain system and brain area. The result is that in any given child, the individual history of maltreatment results in a unique pattern of altered neural systems and resulting psychopathology. The result is that developmental maltreatment is the Great Impostor. Depending up the age, nature and pattern of maltreatment a child may develop symptoms that mimic dozens of traditional DSM-IV diagnoses from autism or ADHD or “learning disorder.” Further, developmental stressors may “express” underlying genetic vulnerabilities; a genetically-vulnerable child may develop a pathological phenotype while a hardier child may not.

This complexity poses a fundamental challenge to any attempts to create simple over-inclusive descriptive categories of psychopathology. Future classification of human psychopathology will need to incorporate a more neurodevelopmentally-informed perspective in order to accurately understand the mechanisms which underlie neuropsychiatric symptoms in a given child or adult. This more mechanism-focused classification would approximate the current model of diagnostic classification in other areas of medicine where there is a more direct connection between the disease process and the pathophysiology. The hope and the promise is that understanding the mechanisms underlying the psychopathology will lead to more effective interventions, and, ultimately, more important, changes in practice, programs and policy that will help prevent the development of abuse-related psychopathology.




Figure 1. Brain organization and monoamine systems: The human brain has a hierarchical organization. The multiple parallel systems in the brain are organized in various brain regions with the most simple in the brainstem and the most complex in the cortex. While somewhat simplified, it is clear that functional complexity correlates with the organizational complexity of the brain, with the most simple regulatory functions mediated by the lower less complex brainstem and the most complex functions - those functions that confer the most unique human properties - are in the cortex. The brainstem and diecephalic originating monoamine systems (DA: dopamine; NE: norepinephrine; 5-HT: serotonin) project up (and down) throughout the brain.




Fig. 2: Impact of neglect on brain development: These images illustrate the impact of neglect on the developing brain. The CT scan on the left is from a healthy three year old child with an average head size (50th percentile). The image on the right is from a three year old child following total global neglect during early childhood. The brain is significantly smaller than average and has abnormal development of cortical, limbic and midbrain structures.

	Event-Related Factors	Individual Characteristics	Family and Social
<u>Increase Risk</u> (Prolong the intensity or duration of the acute stress response)	<ul style="list-style-type: none"> ◆ Multiple or repeated event (in this case, ongoing threat) ◆ Physical injury to child ◆ Involves physical injury or death to loved one, particularly mother ◆ Dismembered or disfigured bodies seen ◆ Destroys home, school or community ◆ Disrupts community infrastructure (as Katrina) ◆ Perpetrator is family member ◆ Long duration, difficult recovery (as in Katrina) 	<ul style="list-style-type: none"> ◆ Female ◆ Age (younger more vulnerable) ◆ Subjective perception of physical harm ◆ History of previous exposure to trauma ◆ No cultural or religious anchors ◆ No shared experience with peers (experiential isolation) ◆ Low IQ ◆ Pre-existing neuropsychiatric disorder (especially anxiety related) 	<ul style="list-style-type: none"> ◆ Trauma directly impacts caregivers ◆ Anxiety in primary caregivers ◆ Continuing threat and disruption to family ◆ Chaotic, overwhelmed family ◆ Physical isolation ◆ Distant caregiving ◆ Absent caregivers
<u>Decrease Risk</u> (Decrease intensity or duration of the acute stress response)	<ul style="list-style-type: none"> ◆ Single event ◆ Perpetrator is stranger ◆ No disruption of family or community structure ◆ Short duration (e.g., tornado) 	<ul style="list-style-type: none"> ◆ Cognitively capable of understanding abstract concepts ◆ Healthy coping skills ◆ Educated about normative post-traumatic responses ◆ Immediate post-traumatic interventions ◆ Strong ties to cultural or religious belief system 	<ul style="list-style-type: none"> ◆ Intact, nurturing family supports ◆ Non-traumatized caregivers ◆ Caregivers educated about normative post-traumatic responses ◆ Strong family beliefs ◆ Mature and attuned parenting skills

Table 1. Risk and attenuating factors for the development of psychopathology following trauma.

- Abercrombie, E. D., & Jacobs, B. L (1987). Single-unit response of noradrenergic neurons in the locus coeruleus of freely moving cats. I. Acutely presented stressful and non-stressful stimuli. *Journal of Neuroscience*, 7, 2837-2847.
- Abercrombie ED, Jacobs BL (1988) Systemic naloxone administration potentiates locus coeruleus noradrenergic neuronal activity under stressful but not non-stressful conditions. *Brain Research* 441:362-366
- Abercrombie ED, Keefe KA, DiFrischia DS, et.al. (1989): Differential effects of stress on in vivo dopamine release in striatum, nucleus accumbens and medial frontal cortex. *Journal of Neurochemistry* 52:1655-1658
- Alexander, P. C., Anderson, C. L., Brand, B., Schaeffer, C. M., Grelling, B. Z., and Kretz, L. (1998) Adult attachment and long-term effects in survivors of incest. *Child Abuse & Neglect* 22[1], 45-61
- Allen, J. R., Heston, J., Durbin, C., and Pruitt, D. B. (1998) Stressors and Development: A Reciprocal Relationship. *Child and Adolescent Psychiatric Clinics of North America* 7[1], 1-18.
- Altman, J., & Das, G.D. (1964). Autoradiographic examination of the effects of enriched environment on the rate of glial multiplication in the adult rat brain. *Nature*, 204, 1161-1165.
- Anda, R. F., Chapman, D. P., Felitti, V. J., Edwards, V. E., Williamson, D. F., Croft, J. P., & Giles, W. H. (2002a). Adverse Childhood Experiences and Risk of Paternity in Teen Pregnancy. *Obstetrics and Gynecology*, 100, 37-45.
- Anda, R. F., Croft, J. B., Felitti, V. J., Nordenberg, D., Giles, W. H., Williamson, D. F., & Giovino, G.A. (1999) Adverse childhood experiences and smoking during adolescence and adulthood. *Journal of the American Medical Association*, 282, 1652.
- Anda, R. F., Felitti, VJ, Chapman D. P., Croft, J. B., Williamson, D. F., Santelli J., Dietz, P. M., & Marks, J. S. (2001) Abused boys, battered mothers, and male involvement in teen pregnancy *Pediatrics*, 107(2)
- Anda, R.F., Felitti, R.F., Walker, J., Whitfield, C., Bremner, D.J., Perry, B.D., Dube, S.R., Giles, W.G. (2006) The enduring effects of childhood abuse and related experiences: a convergence of evidence from neurobiology and epidemiology, *European Archives of Psychiatric and Clinical Neuroscience*, 256 (3) 174 - 186
- Anderson, C. M., Teicher, M. H., Polcari, A., & Renshaw, P. I. (2002) Abnormal T2 relaxation time in the cerebellar vermis of adults sexually abused in childhood: potential role of the vermis in stress-enhanced risk for drug abuse. *Psychoneuroendocrinology*, 27, 231-244.

- Aston-Jones G, Ennis M, Pieribone VA, Nickell WT, Shipley MT (1986), The brain nucleus locus coeruleus: restricted afferent control of a broad efferent network. *Science* 234: 734-7
- Bartanusz V, Jezova D, Bertini LT, Tilders FJ, Aubry JM, Kiss JZ (1993), Stress-induced increase in vasopressin and corticotropin-releasing factor expression in hypophysiotrophic paraventricular neurons. *Endocrinology* 132: 895-902
- Bennett, E.L., Diamond, M.L., Krech, D., & Rosenzweig, M.R. (1964). Chemical and anatomical plasticity of the brain. *Science*, 146, 610-619.
- Bremner JD, Randall P, Vermetten E, et al (1997): Magnetic resonance imaging-based measurement of hippocampal volume in posttraumatic stress disorder related to childhood physical and sexual abuse – a preliminary report. *Biological Psychiatry* 41: 23-32.
- Bennett, A. J, Lesch, K. P., Heils, A., Long, J. C., Lorenz, J. G., Shoaf, S. E., Champoux, M., Suomi, S. J., Linnoila, M. V., & Higley, J. D. (2002). Early experience and serotonin transporter gene variation interact to influence primate CNS function. *Molecular Psychiatry*, 7, 118-122.
- Bookstein, F.L., Streissguth, A.P., Sampson, P.D., Connor, P.D., & Barr, H.M. (2002). Corpus callosum shape and neuropsychological deficits in adult males with heavy fetal alcohol exposure. *NeuroImage*, 15(1), 233-251
- Bremner, J. D. (2003). Long-term effects of childhood abuse on brain and neurobiology. *Child and Adolescent Psychiatric Clinics of North America*, 12, 271-292.
- Bremner, J. D., Narayan, M., Staib, L. H., Southwick, S. M., McGlashan, T., & Charney, D. S. (1999). Neural correlates of memories of childhood sexual abuse in women with and without posttraumatic stress disorder. *American Journal of Psychiatry*, 156, 1787-1795.
- Bremner, J. D., Randall, P.R. Scott, T.M., Capelli, S., Delaney, R., McCarthy, G., & Charney, D. S. (1995). Deficits in short-term memory in adult survivors of childhood abuse. *Psychiatry Research*, 59, 97-107.
- Bremner, J. D., Randall, P., Vermetten, E., Staib, L., Bronen, R. A., Mazure, C., Capelli, S., McCarthy, G., Innis, R. B., & Charney, D. S. (1997) Magnetic image resonance-based measurement of hippocampal volume in posttraumatic stress disorder related to childhood physical and sexual abuse: a preliminary report. *Biological Psychiatry*, 41, 23-32.
- Bremner, J. D., Southwick, M., Darnell, M. A., & Charney, D. S. (1996b). Chronic PTSD in Vietnam combat veterans: Course of illness and substance abuse. *American Journal of Psychiatry*, 153, 369-375.
- Bremner JD, Vythilingam M, Vermetten E, et al (2003): MRI and PET study of deficits in hippocampal structure and function in women with childhood sexual abuse and posttraumatic stress disorder. *American Journal of Psychiatry* 160:924-932.

- Caldji, C., Francis, D., Sharma, S., Plotsky, P. M., & Meaney, M. J. (2000). The effects of early rearing environment on the development of GABA and central benzodiazepine receptor levels and novelty-induced fearfulness in the rat. *Neuropsychopharmacology*, 22, 219-229.
- Caldji, C., Tannenbaum, B., Sharma, S., Francis, D., Plotsky, P. M., & Meaney, M.J. (1998). Maternal care during infancy regulates the development of neural systems mediating the expression of fearfulness in the rat. *Proceedings of the National Academy of Sciences. USA*, 95, 5335-5340.
- Cannon WB (1914), The emergency function of the adrenal medulla in pain and the major emotions. In: *Am J Physiol*, Vol 33, pp 356-372
- Carrion VG, Weems CF, Eliez S, Patwardhan A, Brown W, Ray R, Reiss AL (2001): Attenuation of frontal asymmetry in pediatric posttraumatic stress disorder. *Biological Psychiatry* 50: 943-951.
- Carrion VG, Weems CF, Ray R, Glaser B, Hessel D, Reiss AL (2002a): Diurnal salivary cortisol in pediatric posttraumatic stress disorder. *Biological Psychiatry* 51: 575-582.
- Carrion VG, Weems CF, Ray R, Reiss AL (2002b): Towards an empirical definition of pediatric PTSD: The phenomenology of PTSD symptoms in youth. *Journal of the American Academy of Child & Adolescent Psychiatry* 41: 166-173.
- Carrion, V. G., & Steiner, H. (2000). Trauma and dissociation in delinquent adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 39, 353-359
- Caspi A, McClay J, Moffitt TE, Mill J, Martin J, Craig IW, Taylor A, Poulton R. (2002) Role of genotype in the cycle of violence in maltreated children. *Science*. 2002 Aug 2;297(5582):851-4;
- Caspi, A., Sugden, K., Moffitt, T. E., Taylor, A., Craig, I. W., Harrington, H., McClay, J., Mill, J., Martin, J., Braithwaite, A., & Poulton, R. (2003). Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science*, 301, 386-389.
- Castro-Alamancos MA, Connors BW (1996), Short-term plasticity of a thalamocortical pathway dynamically modulated by behavioral state. *Science* 272: 274-7
- Chugani, H.T., Behen, M.E., Muzik, O., Juhasz, C. Nagy, F. & Chugani, D.C. (2001) Local brain functional activity following early deprivation: a study of post-institutionalized Romanian orphans. *Neuroimage* 14:6 1290-1301.
- Clugnet MC, LeDoux JE (1990), Synaptic plasticity in fear conditioning circuits: induction of LTP in the lateral nucleus of the amygdala by stimulation of the medial geniculate body. *J Neurosci* 10: 2818-24
- Coleman, P.D., & Riesen, A.H. (1968). Environmental effects on cortical dendritic fields: I. rearing in the dark. *Journal of Anatomy (London)*, 102, 363-374.
- Coplan JD, Andrews MW, Rosenblum LA, Owens MJ, Friedman S, Gorman JM, Nemeroff CB (1996), Persistent elevations of cerebrospinal fluid concentrations of corticotropin-releasing factor in adult nonhuman primates exposed to early-life stressors:

- implications for the pathophysiology of mood and anxiety disorders. In: Proc Natl Acad Sci U S A, Vol 93, pp 1619-23
- Cortes A, Saltzman KM, Weems CF, Regnault HP, Reiss AL, Carrion VG (2005): Development of Anxiety Disorders in a Traumatized Pediatric Population: A Longitudinal Evaluation. *Child Abuse and Neglect* 29, 905-914.
- Cummins, R.A., & Livesey, P. (1979). Enrichment-isolation, cortex length, and the rank order effect. *Brain Research*, 178, 88-98.
- Darwin, C. (1868). *The variations of animals and plants under domestication*. London: J. Murray
- Davis M (1992a), The role of the amygdala in fear and anxiety. *Annu Rev Neurosci* 15: 353-75
- Davis M (1992b), The role of the amygdala in fear-potentiated startle: implications for animal models of anxiety. *Trends Pharmacol Sci* 13: 35-41
- De Bellis MD, Hall J, Boring AM, Frustaci K, Moritz G (2001): A pilot study of hippocampal volumes in pediatric maltreatment-related posttraumatic stress disorder. *Biological Psychiatry* 50: 305-309.
- De Bellis, M. D., Baum, A. S., Birmaher, B., Keshavan, M. S., Eccard, C. H., Boring, A. M., Jenkins, F. J., & Ryan, N. D. (1999a). A.E. Bennett Research Award. Developmental traumatology. Part I: Biological stress systems. *Biological Psychiatry*, 15, 45:1259-1270.
- De Bellis, M. D., Baum, A. S., Birmaher, B., & Ryan, N. D. (1997). Urinary catecholamine excretion in childhood overanxious and posttraumatic stress disorders. *Annals of the New York Academy of Science*, 821, 451-455.
- DeBellis, M. D., Chrousos, G. P., Dorn, L. D., Burke L, Helmers, K., Kling, M. A., Trickett, P. K., & Putnam, F. W. (1994). Hypothalamic-pituitary-adrenal axis dysregulation in sexually abused girls. *Journal of Clinical Endocrinology and Metabolism*, 78, 249-255.
- De Bellis, M. D., Keshavan, M. S., Clark, D.B., Casey, B. J., Giedd, J. N., Boring, A. M., Frustaci, K., & Ryan, N. D. (1999b). A.E. Bennett Research Award. Developmental traumatology. Part II: Brain development. *Biological Psychiatry*, 1999, 45, 1271-1284.
- De Bellis, M. D., Keshavan, M. S., Shifflett, H., Iyengar, S., Beers, S. R., Hall, J., & Moritz, G. (2002). Brain structures in pediatric maltreatment-related posttraumatic stress disorder: a sociodemographically matched study. (2002). *Biological Psychiatry*, 52, 1066-78.
- De Bellis, M., & Thomas, L. (2003). Biologic findings of post-traumatic stress disorder and child maltreatment. *Current Psychiatry Reports*, 5, 108-117.
- Dennis, W. (1973). *Children of the Creche*. New York: Appleton-Century-Crofts.

- Diamond, M.C., & Hopson, J. (1998). *Magic Trees of the Mind: How to nurture your child's intelligence, creativity, and healthy emotions from birth through adolescence*. New York: Dutton.
- Diamond, M.C., Krech, D., & Rosenzweig, M.R. (1964). The effects of an enriched environment on the histology of the rat cerebral cortex. *Comparative Neurology*, 123, 111-119.
- Diamond, M.C., Law, F., Rhodes, H., Lindner, B., Rosenzweig, M.R., Krech, D., & Bennett, E.L. (1966). Increases in cortical depth and glia numbers in rats subjected to enriched environments. *Comparative Neurology*, 128, 117-126.
- Dietz, P. M., Spitz, A. M., Anda, R. F., Williamson, D. F., McMahon, P..M., Santelli J. S., Nordenberg, D. F., Felitti, V. J., & Kendrick, J. S. (1999). Unintended pregnancy among adult women exposed to abuse or household dysfunction during their childhood. *Journal of the American Medical Association*, 282, 1359-1364.
- Dong, M., Anda, R. F., Felitti, V. J., Dube, S. R., & Giles, W. H. (2003a). The relationship of exposure to childhood sexual abuse to other forms of abuse, neglect and household dysfunction during childhood. *Child Abuse and Neglect*, 27, 625-639.
- Dong, M., Dube, S. R., Felitti, V. J., Giles, W. H., & Anda, R. F. (2003b). Adverse childhood experiences and self-reported liver disease: new insights into a causal pathway. *Archives of Internal Medicine*, 163, 1949-1956.
- Dube, S. R., Anda, R. F., Felitti, V. J., Chapman, D., Williamson, D. F., & Giles, W. H. (2001a). Childhood abuse, household dysfunction and the risk of attempted suicide throughout the life span: findings from the Adverse Childhood Experiences Study. *Journal of the American Medical Association*, 286, 3089-3096.
- Dube, S. R., Anda, R. F., Felitti, V. J., Croft, J. B., Edwards, V. J., & Giles, W. H. (2001b). Growing up with Parental alcohol abuse: Exposure to childhood abuse, neglect and household dysfunction. *Child Abuse and Neglect*, 25, 1627-1640.
- Dube, S. R., Anda, R. F., Felitti, V. J., Edwards, V. J., & Croft, J. B. (2002a). Adverse childhood experiences and personal alcohol abuse as an adult. *Addictive Behaviors*, 27, 713-725.
- Dube, S. R., Anda, R. F., Felitti, V. J., Edwards, V. J., & Williamson, D. F. (2002b). Exposure to abuse, neglect and household dysfunction among adults who witnessed intimate partner violence as children. *Violence and Victims*, 17, 3-17. -1658.
- Dube, S. R., Felitti, V. J., Chapman, D. P., Giles, W. H., & Anda, R. F. (2003a). Childhood abuse, neglect and household dysfunction and the risk of illicit drug use: the Adverse Childhood Experience Study. *Pediatrics*, 111, 564-572.
- Dube, S. R., Felitti, V. J., Dong, M., Giles, W. H., & Anda, R. F. (2003b). The impact of adverse childhood experiences on health problems: evidence from four birth cohorts dating back to 1900. *Preventive Medicine*, 37, 268-277.

- Dube SR, Williamson DF, Thompson T, Felitti VJ, & Anda RF. (2004). Assessing the Reliability of Retrospective Reports of Adverse Childhood Experiences Among Adult HMO Members Attending a Primary Care Clinic. *Child Abuse and Neglect*, 28(7), 729-737.
- Driessen, M., Herrmann, J., Stahl, K., Zwaan, M., Meier, S., Hill, A., Osterheider, M., & Petersen, D. (2000). Magnetic resonance imaging volumes of the hippocampus and the Amygdala in women with borderline personality disorder and early traumatization. *Archives of General Psychiatry*, 57, 1115-1122.
- Ebinger, P. (1974). A cytoarchitectonic volumetric comparison of brains in wild and domestic sheep. *Z.Anat.Entwicklungsgesch*, 144, 267-302.
- Elam M, Thoren P, Svensson TH (1986), Locus coeruleus neurons and sympathetic nerves: activation by visceral afferents. *Brain Res* 375: 117-25
- Epstein, J. N., Saunders, B. E., Kilpatrick, D. G., and Resnick, H. S. (1998) PTSD as a mediator between childhood rape and alcohol use in adult women. *Child Abuse & Neglect* 22[3], 223-234.
- Felitti, V. J., Anda, R. F., Nordenberg, D, Williamson, D. F., Spitz, A. M., Edwards, V., Koss, M. P., & Marks, J. S. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences Study. *American Journal of Preventive Medicine*, 14, 245-258.
- Foley DL, Eaves LJ, Wormley B, Silberg JL, Maes HH, Kuhn J, Riley B. Childhood adversity, monoamine oxidase a genotype, and risk for conduct disorder. *Arch Gen Psychiatry*. 2004 Jul;61(7):738-44.
- Foote, S. L., Bloom, E., & Aston-Jones, G. (1983). Nucleus locus coeruleus: new evidence of anatomical and physiological specificity. *Physiology and Behavior*, 63, 844-914.
- Francis, D. D., Caldji, C., Champagne, F., Plotsky, P. M, & Meaney, M. J. (1999). The role of corticotropin-releasing factor-norepinephrine systems in mediating the effects of early experience on the development of behavioral and endocrine responses to stress. *Biological Psychiatry*, 46, 1153-1166.
- Francis, D. D., Young, L. J., Meaney, M. J., & Insel, T. R. (2002). Naturally occurring differences in maternal care are associated with the expression of oxytocin and vasopressin (V1a) receptors: gender differences. *Journal of Neuroendocrinology*, 14, 349-353.
- Glaser D. (2000). Child abuse and neglect and the brain: a review. *Journal of Child Psychology and Psychiatry*, 41, 97-116.
- Glavin GB (1985) Stress and brain noradrenaline: a review. *Neuroscience Biobehavioral Review* 9:233-243
- Gorman, J. .M., Mathew, S., & Coplan, J. (2002). Neurobiology of early life stress: nonhuman primate models. *Clinical Neuropsychiatry*, 7, 96-103.
- Gould, E., Reeves, A.J., Graziano, M.S.A., & Gross, C.G. (1999). Neurogenesis in the neocortex of adult primates. *Science*, 286, 548-552.

- Greenough, W.T., & Volkmar, F.R. (1973). Pattern of dendritic branching in occipital cortex of rats reared in complex environments. *Experimental Neurology*, 40, 491-504.
- Greenough, W.T., Volkmar, F.R., & Juraska, J.M. (1973). Effects of rearing complexity on dendritic branching in frontolateral and temporal cortex of the rat. *Experimental Neurology*, 41, 371-378.
- Gunnar, M. R., Morison, S. J., Chisholm, K., & Schuder, M. (2001). Salivary cortisol levels in children adopted from Romanian orphanages. *Development and Psychopathology*, 13(3), 611-628.
- Heffernan, K., Cloitre, M., Tardiff, K., Marzuk, P.M., Portera, L., & Leon, A. C. (2000). Childhood trauma as a correlate of lifetime opiate use in psychiatric patients. *Addictive Behaviors*, 25, 797-803.
- Heidenreich, F.W. (1834). Kaspar Hausers verwundung, krankheit und liechenoffnung. *Journal der Chirurgie und Augen-Heilkunde*, 21 (1834), 91-123.
- Heim C, & Nemeroff, C. B. (2001). The role of childhood trauma in the neurobiology of mood and anxiety disorders: preclinical and clinical studies. *Biological Psychiatry*, 49, 1023-1039.
- Heim, C., Newport, D. J., Bonsall, R., Miller, A.H., & Nemeroff, C. B. (2001). Altered pituitary-adrenal axis responses to provocative challenge tests in adult survivors of childhood abuse. *American Journal of Psychiatry*, 158, 575-81.
- Heim, C., Newport, D. J., Heit, S. Graham, Y. P., Wilcox, M., Bonsall, R., Miller, A. H., & Nemeroff, C. B. (2000). Pituitary-adrenal and autonomic responses to stress in women after sexual and physical abuse in childhood. *Journal of the American Medical Association*, 284, 592-597.
- Heinsbroek RPW, van Haaren F, Feenstra MPG, et.al. (1990) Sex differences in the effects of inescapable footshock on central catecholaminergic and serotonergic activity. *Pharmacol.Biochem.Behav.* 37:539-550
- Heinsbroek RPW, van Haaren F, Feenstra MPG, et.al. (1991) Controllable and uncontrollable footshock and monoaminergic activity in the frontal cortex of male and female rats. *Brain Research* 551:247-255.
- Henry JP, Stephens PM, Ely DL(1986) Psychosocial hypertension and the defence and defeat reactions. *J Hypertension* 4:687-697
- Henry JP, Liu YY, Nadra WE, Qian CG, Mormede P, Lemaire V, Ely D, Hendley ED (1993) Psychosocial stress can induce chronic hypertension in normotensive strains of rats. *Hypertension* 21:714-723
- Herman JP, Guillonneau R, Dantzer R, et.al. (1982) Differential effects of inescapable footshocks and of stimuli previously paired with inescapable footshocks on dopamine turnover in cortical and limbic areas of the rat. *Life Science* 30:2207-2214

- Hubel, D.H., & Wiesel, T.N. (1963). Receptive fields of cells in striate cortex of very young, visually inexperienced kittens. *Journal of Neurophysiology*, 26, 994-1002.
- Hubel, D.H., & Wiesel, T.N. (1970). The period of susceptibility to the physiological effects of unilateral eye closure in kittens. *Journal of Physiology*, 206, 419-436.
- Huttenlocher, J., Vasilyeva, M., Cymerman, E., & Levine, S.C. (2002). Language input and child syntax. *Cognitive Psychology*, 45(3), 337-374.
- Huttenlocher, P.R. (1979). Synaptic density in human frontal cortex: developmental changes and effects of aging. *Brain Research*, 163, 195-205.
- Huttenlocher, P.R. (1994). Synaptogenesis in human cerebral cortex. In G. Dawson & K.W. Fischer (Eds.), *Human Behavior and the Developing Brain*. (pp. 35-54). New York: Guilford.
- Ito, Y., Teicher, M. H., Glod, C. A., & Ackerman, E. (1998) Preliminary evidence for aberrant cortical development in abused children: a quantitative EEG study. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 10, 298-307.
- Kalivas PW (1985) Sensitization to repeated enkephalin administration into the ventral tegmental area of the rat, II: involvement of the mesolimbic dopamine system. *Journal Pharmacolog.Exp.Ther.* 235:544-550
- Kalivas PW, Duffy P, Dilts R, et.al. (1988): Enkephalin modulation of A10 dopamine neurons: a role in dopamine sensitization. *Ann.N.Y.Acad.Sci.* 537:405-414
- Kalivas PW, Richardson-Carlson R, Van Orden G (1986) Cross sensitization between foot shock stress and enkephalin- induced motor activity. *Biological Psychiatry* 21:939-950
- Kehne, John, (2007) The CRF1 Receptor, a Novel Target for the Treatment of Depression, Anxiety, and Stress-Related Disorders: *CNS & Neurological Disorders - Drug Targets*, 2007, 6, 000-000
- Kendler, K. S., Bulik, C. M., Silberg, J., Hettema, J. M., Myers, J., & Prescott, C. A. (2000). Childhood sexual abuse and adult psychiatric and substance abuse disorders in women: an epidemiological and cotwin control analysis. *Archives of General Psychiatry*, 57, 953-959.
- Kluver H, Bucy PC (1937), "Psychic blindness" and other symptoms following bilateral temporal lobectomy in rhesus monkeys. *Am J Physiol* 119: 352-353
- Kuan, C.-Y., Roth, K.A., Flavell, R.A., & Rakic, P. (2000). Mechanisms of programmed cell death in the developing brain. *Trends in Neuroscience*, 23, 291-297.
- Lauder, J.M. (1988). Neurotransmitters as morphogens. *Progress in Brain Research*, 73, 365-388.
- LeDoux JE, Cicchetti P, Xagoraris A, Romanski LM (1990), The lateral amygdaloid nucleus: sensory interface of the amygdala in fear conditioning. *J Neurosci* 10: 1062-9

- LeDoux JE, Iwata J, Cicchetti P, Reis DJ (1988), Different projections of the central amygdaloid nucleus mediate autonomic and behavioral correlates of conditioned fear. *J Neurosci* 8: 2517-29
- Lemieux, A. M., & Coe, C. L. (1995). Abuse-related posttraumatic stress disorder: Evidence for chronic neuroendocrine activation in women. *Psychosomatic Medicine*, 57, 105-115.
- Levine ES, Litto WJ, Jacobs BL (1990), Activity of cat locus coeruleus noradrenergic neurons during the defense reaction. *Brain Res* 531: 189-95
- Levine S, Atha K, Wiener SG (1993a), Early experience effects on the development of fear in the squirrel monkey. In: *Behav Neural Biol*, Vol 60, pp 225-33
- Levine S, Wiener SG, Coe CL (1993b), Temporal and social factors influencing behavioral and hormonal responses to separation in mother and infant squirrel monkeys. In: *Psychoneuroendocrinology*, Vol 18, pp 297-306
- Maunsell JH (1995), The brain's visual world: representation of visual targets in cerebral cortex. In: *Science*, Vol 270, pp 764-9
- McEwen BS (2001), From molecules to mind. Stress, individual differences, and the social environment. In: *Ann N Y Acad Sci*, Vol 935, pp 42-9
- Meaney, M.J., Aitken, D.H., van Berkal, C., Bhatnagar, S., & Sapolsky, R.M. (1988). Effect of neonatal handling on age-related impairments associated with the hippocampus. *Science*, 239 :766-768.
- Miaskowski C, Ong GL, Lukic D, Haldar J (1988), Immobilization stress affects oxytocin and vasopressin levels in hypothalamic and extrahypothalamic sites. *Brain Res* 458: 137-41
- Miczek KA, Thompson ML, Tornatzky W (1990) Subordinate animals: Behavioral and physiological adaptations and opioid tolerance. In *Stress: Neurobiology and Neuroendocrinology*. Edited by MR Brown, GF Koob, C Rivier, New York, Marcel Dekker, pp 323-357
- Molnar, B. E., Shade, S. B., Kral, A. H., Booth, R. E., and Watters, J. K. (1998) Suicidal behavior and sexual/physical abuse among street youth. *Child Abuse & Neglect* 22[3], 213-222
- Money, J., & Anecillo, C. (1976). IQ changes following change of domicile in the syndrome of reversible hyposomatotropinism (psychosocial dwarfism): pilot investigation . *Psychoneuroendocrinology*, 1, 427-429.
- Moore RY, Bloom FE (1979), Central catecholamine neuron systems: anatomy and physiology of the norepinephrine and epinephrine systems. *Annu Rev Neurosci* 2: 113-68
- Morilak DA, Fornal CA, Jacobs BL (1987a), Effects of physiological manipulations on locus coeruleus neuronal activity in freely moving cats. I. Thermoregulatory challenge. *Brain Res* 422: 17-23

- Morilak DA, Fornal CA, Jacobs BL (1987b), Effects of physiological manipulations on locus coeruleus neuronal activity in freely moving cats. II. Cardiovascular challenge. *Brain Res* 422: 24-31
- Morilak DA, Fornal CA, Jacobs BL (1987c), Effects of physiological manipulations on locus coeruleus neuronal activity in freely moving cats. III. Glucoregulatory challenge. *Brain Res* 422: 32-9
- Munk MH, Roelfsema PR, Konig P, Engel AK, Singer W (1996), Role of reticular activation in the modulation of intracortical synchronization. *Science* 272: 271-4
- Nauta WJ, Whitlock DG (1956), Subcortical projections from the temporal neocortex in *Macaca mulatta*. *J Comp Neurol* 106: 183-212
- Papez JW (1937), A proposed mechanism of emotion. *Arch of Neurology and Psychiatry* 38: 725-740
- Pavlidis C, Watanabe Y, McEwen BS (1993a), Effects of glucocorticoids on hippocampal long-term potentiation. In: *Hippocampus*, Vol 3, pp 183-92
- Pavlidis C, Watanabe Y, McEwen BS (1993b), Effects of glucocorticoids on hippocampal long-term potentiation. *Hippocampus* 3: 183-92
- Perry, B.D. (1988). Placental and blood element neurotransmitter receptor regulation in humans: potential models for studying neurochemical mechanisms underlying behavioral teratology. *Progress in Brain Research*, 73, 189-206.
- Perry, B.D. (1994). Neurobiological sequelae of childhood trauma: post-traumatic stress disorders in children. In M. Murberg (Ed.), *Catecholamines in Post-traumatic Stress Disorder: Emerging Concepts*. (pp. 253-276). Washington, D.C.: American Psychiatric Press.
- Perry, B.D. (2001a) The neurodevelopmental impact of violence in childhood. In *Textbook of Child and Adolescent Forensic Psychiatry*, (Eds., D. Schetky and E.P. Benedek) American Psychiatric Press, Inc., Washington, D.C. pp. 221-238
- Perry, B.D. (2001b) The neuroarcheology of childhood maltreatment: the neurodevelopmental costs of adverse childhood events. In: *The Cost of Maltreatment: Who Pays? We All Do*. (Eds., K. Franey, R. Geffner & R. Falconer), Family Violence and Sexual Assault Institute, San Diego, pp. 15-37
- Perry, B.D. (2002) Childhood experience and the expression of genetic potential: what childhood neglect tells us about nature and nurture. *Brain and Mind* 3: 79-100
- Perry, B.D. (2006) *The Neurosequential Model of Therapeutics: Applying principles of neuroscience to clinical work with traumatized and maltreated children* In: *Working with Traumatized Youth in Child Welfare* (Ed. Nancy Boyd Webb), The Guilford Press, New York, NY, pp. 27-52
- Perry, B.D., & Pollard, R. (1997). Altered brain development following global neglect in early childhood. *Proceedings from the Society for Neuroscience Annual Meeting* (New Orleans), (abstract)

- Perry, B.D., & Pollard, R. (1998). Homeostasis, stress, trauma, and adaptation: A neurodevelopmental view of childhood trauma. *Child and Adolescent Psychiatric Clinics of North America*, 7, 33-51.
- Perry, B.D., Pollard, R., Blakely, T., Baker, W., & Vigilante, D. (1995) Childhood trauma, the neurobiology of adaptation and 'use-dependent' development of the brain: How "states" become "traits". *Infant Mental Health J*, 16 (4): 271-291, 1995.
- Perry BD, Stolk JM, Vantini G, Guchhait RB, U'Prichard DC (1983), Strain differences in rat brain epinephrine synthesis: regulation of alpha-adrenergic receptor number by epinephrine. *Science* 221: 1297-9
- Phillips RG, LeDoux JE (1992a), Differential contribution of amygdala and hippocampus to cued and contextual fear conditioning. In: *Behav Neurosci*, Vol 106, pp 274-85
- Phillips RG, LeDoux JE (1992b), Differential contribution of amygdala and hippocampus to cued and contextual fear conditioning. *Behav Neurosci* 106: 274-85
- Plotsky, P.M., & Meaney, M.J. (1993). Early, postnatal experience alters hypothalamic corticotropin releasing factor (CRF) mRNA, median eminence CRF content and stress-induced release in adult rats. *Molec Brain Res*, 18, 195-200.
- Putnam, F. W. (1998). Developmental pathways in sexually abused girls. Presented at *Psychological Trauma: Maturation Processes and Psychotherapeutic interventions*. Harvard Medical School, Boston MA. March 20.
- O'Connor, C., Rutter, M., & English and Romanian Adoptees study team. (2000). Attachment disorder behavior following early severe deprivation: extension and longitudinal follow-up. *J.Am.Acad.Child Adolesc.Psychiatry*, 39, 703-712.
- O'Keefe, M. (1998) Posttraumatic stress disorder among incarcerated battered women: A comparison of battered women who killed their abusers and those incarcerated for other offenses. *Journal of Traumatic Stress* 11[1], 71-85
- Orr, S. P., Lasko, N. B., Metzger, L. J., Berry, N. J., Ahern, C. E., and Pitman, R. K. (1998) Psychophysiological assessment of women with posttraumatic stress disorder resulting from childhood sexual abuse. *Journal of Consulting and Clinical Psychology* 66[6], 906-913
- Rakic, P. (1981). Development of visual centers in the primate brain depends upon binocular competition before birth. *Science*, 214, 928-931.
- Rakic, P. (1996). Development of cerebral cortex in human and non-human primates. In M. Lewis (Ed.), *Child and Adolescent Psychiatry: A Comprehensive Textbook*. (pp. 9-30). New York: Williams and Wilkins.
- Read, J., Perry, B. D., Moskowitz, A., & Connolly, J. (2001). The contribution of early traumatic events to schizophrenia in some patients: a traumagenic neurodevelopmental model. *Psychiatry*, 2001, 64, 319-345

- Rehkamper, G., Haase, E., & Frahm, H.D. (1988). Allometric comparison of brain weight and brain structure volumes in different breeds of the domestic pigeon, *Columba livia* f. d. *Brain Behav.Evol.*, 31, 141-149.
- Reul JM, de Kloet ER (1985), Two receptor systems for corticosterone in rat brain: microdistribution and differential occupation. *Endocrinology* 117: 2505-11
- Rohrs, M. (1955). Vergleichende untersuchungen an wild- und hauskatzen. *Zool.Anz.*, 155, 53-69.
- Rohrs, M., & Ebinger, P. (1978). Die beurteilung von hirngrobenunterschieden zwischen wild- und haustieren. *Z.zool.Syst.Evolut.-forsch*, 16, 1-14.
- Rorty, M. and Yager, J. (1996) Histories of childhood trauma and complex post-traumatic sequelae in women with eating disorders. *The Psychiatric Clinics of North America* 19[4]
- Rosenbaum JF, Biederman J, Gersten M, Hirshfeld DR, Meminger SR, Herman JB, Kagan J, Reznick JS, Snidman N (1988), Behavioral inhibition in children of parents with panic disorder and agoraphobia. A controlled study. *Arch Gen Psychiatry* 45: 463-70
- Rutledge, L.T., Wright, C., & Duncan, J. (1974). Morphological changes in pyramidal cells of mammalian neocortex associated with increased use. *Experimental Neurology*, 44, 209-228.
- Rutter, M., Andersen-Wood, L., Beckett, C., Bredenkamp, D., Castle, J., Grootheus, C., Keppner, J., Keaveny, L., Lord, C., O'Connor, T.G., & English and Romanian Adoptees study team. (1999). Quasi-autistic patterns following severe early global privation. *J.Child Psychol.Psychiat.*, 40, 537-49.
- Rutter, M., & English and Romanian Adoptees study team. (1998). Developmental catch-up, and deficit, following adoption after severe global early privation. *J.Child Psychol.Psychiat.*, 39, 465-476.
- Saper CB (1982), Convergence of autonomic and limbic connections in the insular cortex of the rat. *J Comp Neurol* 210: 163-73
- Sapolsky RM (2000), Glucocorticoids and hippocampal atrophy in neuropsychiatric disorders. In: *Arch Gen Psychiatry*, Vol 57, pp 925-35
- Sapolsky RM, Krey LC, McEwen BS (1984), Glucocorticoid-sensitive hippocampal neurons are involved in terminating the adrenocortical stress response. In: *Proc Natl Acad Sci U S A*, Vol 81, pp 6174-7
- Selden NR, Everitt BJ, Jarrard LE, Robbins TW (1991), Complementary roles for the amygdala and hippocampus in aversive conditioning to explicit and contextual cues. In: *Neuroscience*, Vol 42, pp 335-50
- Schaaf, K. K. and McCanne, T. R. (1998) Relationship of childhood sexual, physical and combined sexual and physical abuse to adult victimization and posttraumatic stress disorder. *Child Abuse & Neglect* 22[11], 1119-1133

- Singer W (1995), Development and plasticity of cortical processing architectures. In: Science, Vol 270, pp 758-64
- Simon, N. (1978) Kaspar Hauser Journal of Autism and Childhood Schizophrenia 8: 209-217.
- Spitz, R.A. (1945). Hospitalism: An inquiry into the genesis of psychiatric conditions in early childhood. Psychoanalytic Study of the Child, 1:53-74.
- Spitz, R.A. (1946). Hospitalism: A follow-up report on investigation described in Volume I, 1945. Psychoanalytic Study of the Child, 2:113-117.
- Strathearn, L.I., Gray, P.H., O'Callaghan, M.J. & Wood, D.W. (submitted) Cognitive neurodevelopment in extremely low birth weight infants: nature vs. nurture revisited
- Stanton ME, Gutierrez YR, Levine S (1988), Maternal deprivation potentiates pituitary-adrenal stress responses in infant rats. In: Behav Neurosci, Vol 102, pp 692-700
- Stein, M. B., Koverola, C., Hanna, C., Torchia, M.G., & McClarty, B (1997). Hippocampal volume in women victimized by childhood sexual abuse. Psychological Medicine, 27, 1-9
- Teicher, M. H., Andersen S.L., Polcari, A, Anderson, C. M., & Navalta, C. P. (2002). Developmental neurobiology of childhood stress and trauma. Psychiatric Clinics of North America, 25, 397-426
- Teicher, M. H., Ito, Y., Glod, C. A., Andersen, S. L., Dumont, N., & Ackerman, E. (1997). Preliminary evidence for abnormal cortical development in physically and sexually abused children using EEG coherence and MRI. Annals of the New York Academy of Sciences, 821, 160-175.
- Teicher, M.H., Ito, Y., Glod, C. A., Schiffer, F., & Ackerman, E. (1994) Possible effects of early abuse on human brain development, as assessed by EEG coherence. American Journal of Neuropsychopharmacology, 33, 52.
- Teicher MH, Dumont NL, Ito Y, Vaituzis C, Giedd JN, Andersen SL. (2004) Childhood neglect is associated with reduced corpus callosum area. Biol Psychiatry 56(2):80-5;
- Thomas LA, De Bellis MD. Pituitary volumes in pediatric maltreatment-related posttraumatic stress disorder. Biol Psychiatry. 2004 Apr 1;55(7):752-8;
- Uno, H., Tarara, R., Else, J., & et.al. (1989). Hippocampal damage associated with prolonged and fatal stress in primates. Journal of Neuroscience, 9, 1705-1711.
- van Cleve, T.C. (1972) The Emperor Frederick II of Hohenstufen, Immutator Mundi Oxford University Press, Oxford pp. 335.
- Veldhuis HD, De Kloet ER (1982), Significance of ACTH4-10 in the control of hippocampal corticosterone receptor capacity of hypophysectomized rats. Neuroendocrinology 34: 374-80
- Wainwright, M.S., Perry, B.D., Won, L.A., O'Malley, K.L., Wang, W-Y & Heller, A. (1995) Immortalization of murine striatal neurons by somatic cell fusion with the N18TG2

neuroblastoma: Characterization of cell lines expressing a variety of dopamine receptors and cholinergic markers. *J Neuroscience*, 15 (1):676-688

Young WS, 3rd, Lightman SL (1992), Chronic stress elevates enkephalin expression in the rat paraventricular and supraoptic nuclei. *Brain Res Mol Brain Res* 13: 111-7