

Shoulder Arthritis

Victoria Division of Family Medicine
"Dine and Learn"

Dave Saliken MD FRCSC

February 19, 2020

Rebalance^{MD}

Objectives

1. Management of Shoulder OA
2. Imaging in Shoulder OA
3. Who gets a shoulder replacement
4. Types of shoulder replacement
5. Post-op Expectations

Rebalance^{MD}

Non-op Management

- Activity Modification
- Physio
- Pain medication
 - Acetaminophen
 - NSAID
 - Avoid opioids if possible
- Injections:
 - Corticosteroid
 - Viscosupplement
 - Do NOT do within 4-6 months of surgery (risk of infection)

Rebalance^{MD}

Who Gets a Shoulder Replacement?

- Significant pain and/or shoulder dysfunction associated with glenohumeral cartilage wear
- Have trialed a structured non-operative management protocol
- Typical age >60
- But some young patients: recurrent dislocators, fracture, AVN


Rebalance^{MD}

Imaging Work-up

- Xray
 - Bone loss, cuff
- Surgeon Might Order:
 - U/S or MRI
 - If concerned about rotator cuff integrity
 - CT
 - For surgical planning/templating

Rebalance^{MD}

Primary GH OA



Imaging - Xray

Normal Shoulder Xray




Cuff Tear Arthropathy




Rebalance^{MD}

CT Imaging – Glenoid Anatomy




Rebalance^{MD}

Bone Stock



Walch Classification



Rebalance^{MD}

CT for Surgical Templating



Rebalance^{MD}

Types of Shoulder Arthroplasty

Normal Cuff
Adequate Bone Stock



Anatomic TSA or Hemi



Deficient Cuff
Poor Bone Stock



Reverse TSA



Also more
common in
RA, fracture
sequelae

Rebalance^{MD}

Poor Rotator Cuff → Early Failure of *anatomic* TSA!

- If cuff is deficient or tears we see rapid glenoid loosening with “anatomic” TSA
- In these situations we do a “reverse” TSA instead





Fig. 10. The vertical “rocking horse effect” on the glenoid: ascendant shearing forces leading to loosening of the glenoid component (arrows)


Don't want rocking horse!

Rebalance^{MD}

Reverse TSA - Biomechanics


1) Increased Deltoid
Tension

Lowered center of rotation (COR)



2) More Deltoid
Fibers Engaged

Medialized COR





Rebalance^{MD}

Modular Implants

Shoulder Anatomy is VARIABLE

So need MODULAR IMPLANTS



Rebalance^{MD}

Shoulder Immobilizer

Simple Sling



Neutral Rotation Sling



Abduction Sling



• Baumgarten et al. JSES (2018) 27, 2120–2128

- Improvement in ROM (ER and adduction), less pain with neutral rotation sling

Rebalance^{MD}

Post-op Recovery

- **Weeks 0-6 in sling:**
 - Rx for dilaudid +/- gabapentin
 - Should be off by week 2
 - Exercises (3-5x/day):
 - Passive elbow/hand/wrist ROM
 - Pendulum shoulder
 - Avoid ER beyond neutral
- **Weeks 6-12:**
 - Remove sling
 - Physio begins
 - Focus on ROM
- **Weeks 12-24:**
 - Focus on re-training shoulder function
- **Full recovery typically expected around 6-9 months post-op**
- **Shoulder X-rays: 2 weeks, 6 weeks, 3 months, 6 months then yearly**
 - Ensure no complications

Rebalance^{MD}